

09422999

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1649JXM

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2	Sep 29	The Philippines Inventory of Chemicals and Chemical Substances (PICCS) has been added to CHEMLIST
NEWS	3	Oct 27	New Extraction Code PAX now available in Derwent Files
NEWS	4	Oct 27	SET ABBREVIATIONS and SET PLURALS extended in Derwent World Patents Index files
NEWS	5	Oct 27	Patent Assignee Code Dictionary now available in Derwent Patent Files
NEWS	6	Oct 27	Plasdoc Key Serials Dictionary and Echoing added to Derwent Subscriber Files WPIDS and WPIX
NEWS	7	Nov 29	Derwent announces further increase in updates for DWPI
NEWS	8	Dec 5	French Multi-Disciplinary Database PASCAL Now on STN
NEWS	9	Dec 5	Trademarks on STN - New DEMAS and EUMAS Files
NEWS	10	Dec 15	2001 STN Pricing
NEWS	11	Dec 17	Merged CEABA-VTB for chemical engineering and biotechnology
NEWS	12	Dec 17	Corrosion Abstracts on STN
NEWS	13	Dec 17	SYNTHLINE from Prous Science now available on STN
NEWS	14	Dec 17	The CA Lexicon available in the CAPLUS and CA files
NEWS	15	Jan 05	AIDSLINE is being removed from STN
NEWS	16	Feb 06	Engineering Information Encompass files have new names
NEWS EXPRESS			FREE UPGRADE 5.0e FOR STN EXPRESS 5.0 WITH DISCOVER! (WINDOWS) NOW AVAILABLE
NEWS HOURS			STN Operating Hours Plus Help Desk Availability
NEWS INTER			General Internet Information
NEWS LOGIN			Welcome Banner and News Items
NEWS PHONE			Direct Dial and Telecommunication Network Access to STN
NEWS WWW			CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 09:22:54 ON 15 FEB 2001

=> file medline biosis embase caplus uspatfull

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'MEDLINE' ENTERED AT 09:23:09 ON 15 FEB 2001

FILE 'BIOSIS' ENTERED AT 09:23:09 ON 15 FEB 2001

COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'EMBASE' ENTERED AT 09:23:09 ON 15 FEB 2001

COPYRIGHT (C) 2001 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 09:23:09 ON 15 FEB 2001

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'USPATFULL' ENTERED AT 09:23:09 ON 15 FEB 2001

CA INDEXING COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

=> s camp (a) gef

L1 19 CAMP (A) GEF

=> dup rem l1

PROCESSING COMPLETED FOR L1

L2 8 DUP REM L1 (11 DUPLICATES REMOVED)

=> d l2 total ibib kwic

L2 ANSWER 1 OF 8 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2001:95767 CAPLUS

TITLE: New signaling pathways for hormones and cyclic adenosine 3',5'-monophosphate action in endocrine cells

AUTHOR(S): Richards, JoAnne S.

CORPORATE SOURCE: Department of Molecular and Cellular Biology, Baylor College of Medicine, Houston, TX, 77030, USA

SOURCE: Mol. Endocrinol. (2001), 15(2), 209-218

CODEN: MOENEN; ISSN: 0888-8809

PUBLISHER: Endocrine Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The glycoprotein hormones, ACTH, TSH, FSH, and LH regulate diverse functions in endocrine cells. Although cAMP and PKA have long been shown to mediate specific intracellular signaling events including the transcription of specific genes via the CREB-CBP complex, recent observations have indicated that PKA does not account for all of the intracellular targets of cAMP. For example, TSH stimulation of thyroid cell proliferation is not completely blocked by PKA inhibitors. TSH and FSH can stimulate PKB phosphorylation by a PKA-independent but PI3-K/PDK1-dependent pathway. An FSH inducible kinase, Sgk, has recently been shown to be a close relative of PKB. Sgk is also a target of PI3-K/PDK1 pathway, indicating that some effects previously ascribed to PKB may be mediated by this inducible kinase. The identification of novel

cAMP-binding proteins that exhibit guanine nucleotide exchange (GEF) activity (**cAMP-GEFS**; Epacs) has open new doors for cAMP action that include activation of small GTPases such as Rap1a, Rap2, and possibly Ras. These GTPases are known activators of down-stream kinase cascades, including p38MAPK and Erk1/2 as well as PI3-K. Thus, FSH and TSH activation of PKB and Sgk may occur via this alternative cAMP

pathway that involves **cAMP-GEFs** and the activation of the PI3-K/PDK1 pathway.

L2 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2000:291074 CAPLUS

DOCUMENT NUMBER: 132:318614

TITLE: Mammalian Rap1A and Ras-associated guanine nucleotide exchange proteins and cDNAs and methods for

diagnosis,

therapy, and drug screening

INVENTOR(S): Kawasaki, Hiroaki; Graybiel, Ann; Housman, David

PATENT ASSIGNEE(S): Massachusetts Institute of Technology, USA

SOURCE: PCT Int. Appl., 128 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000024768	A2	20000504	WO 1999-US24826	19991022
WO 2000024768	A3	20001109		

W: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 1998-105507 19981023

US 1998-108685 19981116

AB The present invention describes the identification, isolation, sequencing and characterization of two human CalDAG-GEF, and two human **cAMP-GEF** genes, which are assocd. with the Ras pathway. Also identified are CalDAG-GEF gene homologs in mice and **cAMP-GEF** gene homologs in rats. Nucleic acids and proteins comprising or derived from the CalDAG-GEFs and/or **cAMP-GEFs** are useful in screening and diagnosing certain Ras-assocd. cancers, in identifying and developing therapeutics for treatment of certain Ras-assocd. cancers, and in producing cell lines and transgenic animals useful as models of Ras-assocd. cancers. Thus, CalDAG-GEFI was found to increase GTP-bound Rap1A and this increase was augmented in the presence of the calcium ionophore A23187 or the phorbol ester TPA. CalDAG-GEFI reduced RasV12 activation of Elk1 by .apprx.4-fold. Northern anal. indicated that human CalDAG-GEFI is expressed strongly in the brain and that the mRNA for this protein is strikingly enriched in the striatum. Further, the CalDAG-GEFI is synthesized in striatal projection neurons

and

is transported to striatopallidal and striatonigral terminals.

CalDAG-GEFII activated Ras, but not RalA or Rap1A. Unlike CalDAG-GEFI, CalDAG-GEFII increased the transcriptional activity of Elk1 downstream to Erk/MAP kinase. Northern anal. showed highest expression of CalDAG-GEFII in the cerebellum, cerebral cortex, and amygdala, with low expression in the striatum. **cAMP-GEFI** and **II** strongly and selectively activated Rap1A, but not Ras or RalA, in the presence of **cAMP**. Northern anal. indicated that **cAMP-GEFI** was widely expressed while **cAMP-GEFII** was expressed selectively in the brain and adrenal glands.

IT cDNA sequences

(for mammalian Rap1A and Ras-assocd. guanine nucleotide exchange proteins CalDAG-GEF and **cAMP-GEF**)

IT Protein sequences

(of mammalian Rap1A and Ras-assocd. guanine nucleotide exchange proteins CalDAG-GEF and **cAMP-GEF**)

L2 ANSWER 3 OF 8 MEDLINE DUPLICATE 1
 ACCESSION NUMBER: 2000414769 MEDLINE
 DOCUMENT NUMBER: 20372739
 TITLE: Rap2 as a slowly responding molecular switch in the Rap1 signaling cascade.
 AUTHOR: Ohba Y; Mochizuki N; Matsuo K; Yamashita S; Nakaya M; Hashimoto Y; Hamaguchi M; Kurata T; Nagashima K; Matsuda M
 CORPORATE SOURCE: Department of Pathology, Research Institute, International Medical Center of Japan, Shinjuku-ku, Tokyo 162-8655, Japan.
 SOURCE: MOLECULAR AND CELLULAR BIOLOGY, (2000 Aug) 20 (16) 6074-83.
 Journal code: NGY. ISSN: 0270-7306.
 PUB. COUNTRY: United States
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200011
 ENTRY WEEK: 20001101
 AB . . . 50% of total Rap2 protein in adherent cells. Guanine nucleotide exchange factors (GEFs) for Rap1, C3G, Epac (or cyclic AMP [**cAMP**] -**GEF**), CalDAG-GEFI, PDZ-GEF1, and GFR efficiently increased the level of GTP-Rap2 both in 293T cells and in vitro. GTPase-activating proteins (GAPs). . .

L2 ANSWER 4 OF 8 MEDLINE DUPLICATE 2
 ACCESSION NUMBER: 2000431014 MEDLINE
 DOCUMENT NUMBER: 20420809
 TITLE: Quantitative determination of Rap 1 activation in cyclic nucleotide-treated HL-60 leukemic cells: lack of Rap 1 activation in variant cells.
 AUTHOR: von Lintig F C; Pilz R B; Boss G R
 CORPORATE SOURCE: Department of Medicine and Cancer Center, University of California, San Diego, La Jolla, California, CA 92093-0652,
 USA.
 CONTRACT NUMBER: R01GM055586 (NIGMS)
 R21 CA81115 (NCI)
 SOURCE: ONCOGENE, (2000 Aug 17) 19 (35) 4029-34.
 Journal code: ONC. ISSN: 0950-9232.
 PUB. COUNTRY: ENGLAND: United Kingdom
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals; Cancer Journals
 ENTRY MONTH: 200011
 ENTRY WEEK: 20001103
 AB . . . parental cells. Thus, **cAMP** activation of Rap 1 in HL-60 cells is likely through a **cAMP**-regulated guanine nucleotide exchange factor (**cAMP-GEF**) and since **cAMP** does not activate Rap 1 in the variant cells, the data suggest that full post-translational processing of Rap 1 is necessary for **cAMP-GEF** activation of Rap 1. Activation of Rap 1 by cGMP analogs has not been previously found and suggests possible cross-talk. . .

L2 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2001 ACS
 ACCESSION NUMBER: 2000:721365 CAPLUS
 DOCUMENT NUMBER: 133:317777
 TITLE: Follicle-stimulating hormone (FSH) stimulates phosphorylation and activation of protein kinase B (PKB/Akt) and serum and glucocorticoid-induced kinase (Sgk): evidence for A kinase-independent signaling by FSH in granulosa cells
 AUTHOR(S): Gonzalez-Robayna, Ignacio J.; Falender, Allison E.; Ochsner, Scott; Firestone, Gary L.; Richards, JoAnne

S.
 CORPORATE SOURCE: Department of Molecular and Cellular Biology, Baylor
 College of Medicine, Houston, TX, 77030, USA
 SOURCE: Mol. Endocrinol. (2000), 14(8), 1283-1300
 CODEN: MOENEN; ISSN: 0888-8809
 PUBLISHER: Endocrine Society
 DOCUMENT TYPE: Journal
 LANGUAGE: English
 REFERENCE COUNT: 76
 REFERENCE(S): (1) Adams, S; Nature 1991, V349, P694 CAPLUS
 (2) Adashi, E; Endocrinology 1988, V122, P1583 CAPLUS
 (3) Agati, J; J Biol Chem 1998, V273, P18751 CAPLUS
 (4) Alessi, D; EMBO J 1996, V15, P6541 CAPLUS
 (5) Alliston, T; Endocrinology 2000, V141, P385

CAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

AB FSH stimulates in ovarian granulosa cells diverse, differentiation-
 dependent responses that implicate activation of specific cellular
 signaling cascades. In these studies three kinases were investigated to
 det. their relationship to FSH, cAMP, and A kinase signaling: protein
 kinase B (PKB/Akt), serum and glucocorticoid-induced kinase (Sgk), and
 p38 mitogen-activated protein kinase (p38MAPK). The phosphorylation
 (activation) of these kinases was analyzed by using selective
 agonists/inhibitors: forskolin/H 89 for cAMP-dependent protein kinase (A
 kinase), insulin-like growth factor I (IGF-I)/LY 294002 and wortmannin
 for phosphatidylinositol-dependent kinase (PI3-K), and phorbol myristate
 (PMA)/GF 109203X for diacylglycerol and Ca++-dependent kinases (C
 kinases). An inhibitor (PD 98059) of MEK1, which regulates extracellular
 regulated kinases (ERKs), and SB 203580, which inhibits p38MAPK, were
 also used. In addn., we analyzed the expression of the recently described,
 cAMP-regulated guanine nucleotide exchange factors (cAMP-GEFI and GEFII)
 that impact Ras-related GTPases and Raf kinases, known regulators of
 various protein kinase cascades. We provide evidence that FSH,
 forskolin,
 and 8-bromo-cAMP stimulate phosphorylation of PKB by mechanisms involving
 PI3-K (LY 294002/wortmannin sensitive) not A kinase (H 89 insensitive), a
 pattern of response mimicking that of IGF-I. In contrast, FSH induction
 and phosphorylation of Sgk protein requires A kinase (H 89 sensitive) but
 also involves PI3-K (LY 294002 sensitive) as well as p38MAPK (SB 203580
 sensitive) pathways. PMA (C kinase) abolished FSH-mediated (but not
 IGF-I-mediated) phosphorylation of PKB at a step(s) upstream of PI3-K and
 independent of A kinase. Lastly, FSH-mediated phosphorylation of p38MAPK
 is neg. affected by A kinase and PI3-K, suggesting that it may be
 downstream of specific members of the cAMP-GEF/Rap/Raf
 pathway. We propose that cAMP activation of A kinase is obligatory for
 transcription of Sgk in granulosa cells whereas cAMP
 (IGF-I-like)-mediated phosphorylation (activation) of PKB and Sgk (via PI3-K), as well as
 p38MAPK, involves other cellular events. These results provide new and
 exciting evidence that cAMP acts in granulosa cells by A kinase-dependent
 and -independent mechanisms, each of which controls specific kinase
 cascades.

L2 ANSWER 6 OF 8 BIOSIS COPYRIGHT 2001 BIOSIS DUPLICATE 3
 ACCESSION NUMBER: 2001:27587 BIOSIS
 DOCUMENT NUMBER: PREV200100027587
 TITLE: Expression of cAMP-regulated guanine nucleotide exchange
 factors in pancreatic beta-cells.
 AUTHOR(S): Leech, Colin A.; Holz, George G.; Chepurny, Oleg; Habener,
 Joel F. (1)
 CORPORATE SOURCE: (1) Laboratory of Molecular Endocrinology, Massachusetts
 General Hospital, 55 Fruit Street, WEL320, Boston, MA,
 02114: jhabener@partners.org USA

SOURCE: Biochemical and Biophysical Research Communications,
(November 11, 2000) Vol. 278, No. 1, pp. 44-47. print.
ISSN: 0006-291X.

DOCUMENT TYPE: Article

LANGUAGE: English

SUMMARY LANGUAGE: English

AB. . . remains unknown. Here we present evidence for the expression of
type 1 and type 2 cAMP-regulated guanine nucleotide exchange factors (
cAMP-GEFs) in beta-cells. GEFs are activated by their
binding of cAMP. Because **cAMP-GEFs** activate Ras/MAPK
proliferation signaling pathways, they may play an important role in
PKA-independent, GLP-1-mediated, signaling pathways in the regulation of.

L2 ANSWER 7 OF 8 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1999:136940 CAPLUS

DOCUMENT NUMBER: 130:178854

TITLE: Isolation and characterization of a novel family of
guanine nucleotide exchange factors (GEF) directly
regulated by second messenger systems. Novel second
messenger pathways

AUTHOR(S): Kawasaki, Hiroaki

CORPORATE SOURCE: Cent. Cancer Res., Massachusetts Inst. Technol., USA

SOURCE: Jikken Igaku (1999), 17(4), 490-494

CODEN: JIIGEF; ISSN: 0288-5514

PUBLISHER: Yodosha

DOCUMENT TYPE: Journal; General Review

LANGUAGE: Japanese

AB A review with 12 refs., on isolation, distribution, and the structure of
a

novel family of guanine nucleotide exchange factors (GEF) having second
messenger-binding sites, i.e. **cAMP-GEF** (**cAMP**
-regulated GEF) and CalDAG-GEF (calcium and diacylglycerol-regulated
GEF),
regulation of **cAMP-GEF** and CalDAG-GEF by second
messenger mols., their substrate specificity, and substrate-specific
activation of Rap1 by second messenger mols. through activation of GEF
domain.

L2 ANSWER 8 OF 8 MEDLINE DUPLICATE 4

ACCESSION NUMBER: 1999074384 MEDLINE

DOCUMENT NUMBER: 99074384

TITLE: A family of cAMP-binding proteins that directly activate
Rap1.

AUTHOR: Kawasaki H; Springett G M; Mochizuki N; Toki S; Nakaya M;
Matsuda M; Housman D E; Graybiel A M

CORPORATE SOURCE: Department of Brain and Cognitive Sciences, Massachusetts
Institute of Technology (MIT), Cambridge, MA, 02139, USA.

CONTRACT NUMBER: R01 HD28341 (NICHD)
P01 CA42063 (NCI)
P01 HL41484 (NHLBI)
+

SOURCE: SCIENCE, (1998 Dec 18) 282 (5397) 2275-9.
Journal code: UJ7. ISSN: 0036-8075.

PUB. COUNTRY: United States

LANGUAGE: English

FILE SEGMENT: Priority Journals; Cancer Journals

OTHER SOURCE: GENBANK-U78167; GENBANK-U78168; GENBANK-U78516;
GENBANK-U78517

ENTRY MONTH: 199903

AB . . . brain and body organs and that exhibit both cAMP-binding and
guanine nucleotide exchange factor (GEF) domains is reported. These
cAMP-regulated **GEFs** (**cAMP-GEFs**) bind cAMP
and selectively activate the Ras superfamily guanine nucleotide binding

protein Rap1A in a cAMP-dependent but PKA-independent manner. Our. . .

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

18.63

18.84

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-2.35

-2.35

STN INTERNATIONAL LOGOFF AT 09:24:39 ON 15 FEB 2001